



**PREP INSTRUCTIONS  
FOR COLONOSCOPY  
WITH LYTELY/PEG 3350**

**YOUR COLONOSCOPY IS SCHEDULED FOR:**

**Date of Procedure:** \_\_\_\_\_

**Arrival Time:** \_\_\_\_\_

**Provider:** \_\_\_\_\_

**Facility Name/Address:** \_\_\_\_\_

**Phone:** \_\_\_\_\_

Missed appointments or appointments that are cancelled without a 72 hour notice, may be subject to a fee.

**PLEASE REVIEW THE INSTRUCTIONS AT LEAST 7 DAYS PRIOR TO THE PROCEDURE.**

**USE THESE PREPARATION INSTRUCTIONS. DO NOT USE THE INSTRUCTIONS INCLUDED WITH THE GOLYTELY, GAVILYTE OR PEG 3350.**



**IT IS IMPORTANT THAT YOU FOLLOW THESE DIRECTIONS CAREFULLY AND COMPLETE THE ENTIRE PREPARATION TO ENSURE THE MOST EFFECTIVE CLEANSING.**

- If you are taking blood thinners, insulin or oral medications for diabetes, please let our office know and call the prescribing physician for special dosing instructions.
- Please make sure that you have a responsible adult with you upon arrival at the facility. This adult must stay at the facility during the procedure and drive you home afterwards. You will be unable to drive due to the sedation you will receive. You may not take a Lyft, Uber or other public transportation to or from the facility.

## **7 DAYS PRIOR TO PROCEDURE**

- Stop taking oral Iron supplements. Multivitamins are ok to continue

## **3 DAYS PRIOR TO PROCEDURE**

- Stop eating raw fruits, vegetables, nuts, seeds and beans

## **1 DAY PRIOR TO PROCEDURE**

- You may have clear liquids only
  - Examples include: water, black coffee (sugar is ok), tea, broth, soda, apple juice, Gatorade, Kool-Aid, Jell-O and popsicles
- DO NOT CONSUME ANY DAIRY PRODUCTS OR ANYTHING THAT IS RED, ORANGE, PURPLE OR BLUE
- It is important to drink additional clear liquids before starting and your prep, as well as during, to maintain adequate hydration

## **BEGINNING THE GOLYTELY, GAVILYTE OR PEG 3350**

### **Step 1:**

- For best results mix ahead of time and refrigerate the solution prior to drinking
- If desired, add one of the flavor packets into the container
- Fill the provider container with up to the fill line with lukewarm water
- Shake well (until all powder is dissolved) and place in the refrigerator

### **Step 2:**

- Starting at 5pm, drink one 8 oz. glass every 10-15 minutes until half of the container is gone

**It is essential to drink the amount of liquid instructed. If you experience nausea, bloating or cramping, pause or slow the rate of drinking the additional water until symptoms diminish.**

## **THE DAY OF THE PROCEDURE**

### **Step 4:**

- 6 hours before your arrival time, drink one 8oz glass every 10-15 minutes until the container is empty

**It is essential to drink the amount of liquid instructed. If you experience nausea, bloating or cramping, pause or slow the rate of drinking the additional water until symptoms diminish.**

**YOU MUST COMPLETE STEP 6 AND STOP ALL LIQUIDS 4 HOURS PRIOR TO YOUR ARRIVAL.**